

URSA MAJOR: A MISSING LOW-MASS CDM HALO?

JAN T. KLEYN¹, MARK I. WILKINSON², N. WYN EVANS², GERARD GILMORE²

¹Institute for Astronomy, 2680 Woodlawn Drive, Honolulu, HI, 96822, USA

²Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, UK

Draft version February 5, 2008

ABSTRACT

The recently discovered Ursa Major dwarf spheroidal (dSph) galaxy candidate is about five to eight times less luminous than the faintest previously known dSphs And IX, Draco, and UMi. In this Letter, we present velocity measurements of seven color-magnitude selected UMajor candidate stars. Two of them are apparent non-members based on metallicity and velocity, and the remaining five stars yield a systemic heliocentric velocity of $\bar{v} = -52.45 \pm 4.27$ km s⁻¹ and a central line of sight velocity dispersion of $\langle v^2 \rangle^{1/2} = 9.3^{+11.7}_{-1.2}$ km s⁻¹, with 95% confidence that $\langle v^2 \rangle^{1/2} > 6.5$ km s⁻¹. Assuming that UMajor is in dynamical equilibrium, it is clearly dark matter dominated, and cannot be a purely stellar system like a globular cluster. It has an inferred central mass-to-light ratio of $M/L \sim 500 M_{\odot}/L_{\odot}$ and, based on our studies of other dSphs, may possess a much larger total mass to light ratio. UMajor is unexpectedly massive for its low luminosity – indeed, UMajor appears to be the most dark-matter dominated galaxy yet discovered. The presence of so much dark matter in UMajor immediately suggests that it may be a member of the missing population of low-mass galaxies predicted by the Cold Dark Matter (CDM) paradigm. Given the weak correlation between dSph mass and luminosity, it is entirely likely that a population of dark dwarfs surrounds our Galaxy.

Subject headings: galaxies: individual: Ursa Major dwarf – galaxies: kinematics and dynamics – Local Group – dark matter – celestial mechanics, stellar dynamics

1. INTRODUCTION

All of the Local Group dwarf spheroidal (dSph) galaxies have velocity dispersions much larger than expected for self-gravitating stellar systems, implying that their dynamics is dominated by dark matter, with the stars being little more than dynamical tracers within a dark halo (*e.g.* Mateo 1998). Some dSphs have central mass to light ratios of $\sim 100 M_{\odot}/L_{\odot}$, and average mass to light ratios of several hundred (*e.g.* Kleyna et al. 2001, 2004; Wilkinson et al. 2004).

Recently, Willman et al. (2005) discovered a new candidate dSph in Ursa Major in a search of data from the Sloan Digital Sky Survey (SDSS) (Abazajian et al. 2005). Located about 100 kpc from the Galaxy, its half-light radius is $r_{1/2} \sim 250$ pc, covering $7.75'$ on the sky. Most remarkably, with $M_V \sim -6.75$, it is about five times less luminous than the faintest previously known dSph And IX (Zucker et al. 2004; Chapman et al. 2005) and about eight times less luminous than the faintest Galactic dSphs UMi and Draco (Mateo 1998).

In this paper, we describe precise radial velocity measurements of seven UMajor candidate stars. We compute UMajor's inferred central density and mass, and discuss how UMajor fits into the rest of the rest of the dSph population within the context of standard CDM theory.

2. DATA

We selected candidate stars from Data Release Three of SDSS, picking targets within $6'$ of UMajor's position as given by Willman et al. (2005). Figure 1 shows the color magnitude diagram of the central $6'$ region, slightly smaller than Willman's half-light radius $r_{1/2} = 7.75'$. The giant branch and horizontal branch are both visible, al-

though together they contain only about 50 stars. Our seven targets (dark circles in Figure 1) were drawn from the brightest part of the giant branch, and span the magnitude range $i = 17.45$ to $i = 18.38$. Additionally, we observed the bright velocity standards HD107328, HD90861, and HD132737.

We observed our stars using the upgraded HIRES (Vogt 1994) echelle spectrograph on the Keck I telescope on the night of May 17, 2005. Each star was observed in one integration lasting 1800s. The long spectral coverage of the spectrograph allowed us to obtain wavelengths from H α (6564Å) to the red-most line of the Calcium triplet (8662Å). The signal to noise varied among the spectra from $S/N = 12$ to 3 per pixel, or $S/N = 61$ to 17 per Å. Only the red-most Ca triplet line was near a sky line, but sky subtraction was generally clean even in this case.

We extracted the spectra using the MAKEE data reduction package for HIRES, creating flux and variance spectra for each echelle order. Because the latest version of MAKEE for the new three-chip upgrade to HIRES does not at present solve for the dispersion, we fit the dispersion solution manually for the relevant echelle orders using IRAF. All calibration arc exposures had to be taken during the afternoon or morning because of persistent ghosting with the new chips. To verify instrument stability, we measured the positions of sky lines in the science exposures, and found them to be constant within 0.2 km s⁻¹.

For each of the four stellar absorption lines of interest (H α and the three Ca lines), we cross-correlated a synthetic Gaussian template with the appropriate echelle order using the IRAF FXCOR package. In all cases except for the third Ca triplet line for the faintest UMajor star, we obtained a clear cross-correlation function peak.

A potential problem with single-slit observations is that

mis-centering of the star on the slit will produce a velocity offset common to all orders. To address this problem, we note that telluric absorption lines experience the same spurious velocity shift as stellar absorption lines, and can be used to adjust the velocity back to its correct value. Accordingly, we computed a velocity adjustment using telluric absorption features around 6880Å. Using a template created from the telluric lines from one of the bright standard stars, we used cross-correlation to compute relative velocity adjustments of the other stars. In all cases, the adjustment was less than 0.8 km s^{-1} .

An additional potential source of inaccuracy is template mismatch; for instance, the H α line is actually a blend of several different transitions, and may not be exactly at the template wavelength. To address this problem, we adjusted all of the velocities for each stellar line en masse relative to the second Ca line by computing the median velocity difference between the line and the second Ca line. This procedure introduced a shift of at most 1.6 km s^{-1} .

Next, we treated each velocity for each stellar line as an independent measurement, and combined them to obtain a final velocity and error. Finally, we shifted the entire velocity set to bring the standards into agreement with their published velocities, with a final scatter of 0.4 km s^{-1} , slightly larger than the intrinsic standard star uncertainty of 0.3 km s^{-1} . For each star, we obtained two error estimates: σ_{IRAF} , obtained by rescaling the nominal IRAF FXCOR velocity errors to give the expected χ^2 ; and σ_{scat} , obtained from the empirical velocity scatter among individual lines. For UMajor stars, the two errors are generally similar, but σ_{IRAF} apparently overestimates the errors for the bright standard stars. Table 1 shows the results for our seven target stars.

In Figure 2, we show the velocities as a function of magnitude. Objects 3 and 4 are clearly outliers, separated about 60 km s^{-1} from the other five objects.

The equivalent width of the Ca triplet can be used as a measure of a star's metallicity, for known surface gravity (dwarf vs. giant) (*e.g.* Armandroff & Da Costa 1991). To establish whether the kinematical outliers are members of the UMajor population, we computed the equivalent width of the Ca triplet lines by fitting them with a Lorentzian profile plus a linear continuum, and then integrating the difference between the line profile fit and the continuum fit. We compute the uncertainties of the equivalent widths using a Monte-Carlo procedure: we generated simulated data using our best fit solution added to the empirical noise spectrum output by the MAKEE reduction package, and fit the simulated data in the same manner as the real data.

Figure 3 shows the equivalent widths for the three Ca lines. The kinematical outliers 3 and 4 clearly have a larger equivalent width for the second and third Ca lines, although the first line shows no difference. Object 7 produces very poor fits because the lines are barely discernible above the noise, and its equivalent widths are highly suspect. Generally, only the second and third lines are used for metallicity determination (Armandroff & Da Costa 1991), so the absence of an effect for the weaker first line is not a source of concern. However, for a fixed metallicity, the equivalent width is anti-correlated with magnitude with a slope about $0.6\text{\AA}$ per magnitude (Armandroff & Da

Costa 1991), so that it appears anomalous that star 6 has a larger equivalent width than the brighter stars 1, 2, and 5. To be cautious, we consider the possibility that both 6 and 7 are non-members.

3. VELOCITY DISPERSION AND SYSTEMIC VELOCITY

The simple RMS line of sight dispersion of the five member objects (1,2,5,6, and 7) is 9.1 km s^{-1} , to which measurement errors contribute only slightly. To compute the dispersion $\langle v^2 \rangle^{1/2}$ more rigorously, we assume that the objects are drawn from a Gaussian distribution, and apply equation 2 of Klyna et al. (2004), imposing an *a priori* maximum value of 50 km s^{-1} on the dispersion, and using σ_{scat} as the velocity uncertainty. We do not consider the effect of binary stars, which is generally small. Including all member stars, we obtain $\langle v^2 \rangle^{1/2} = 9.3^{+11.7}_{-1.2} \text{ km s}^{-1}$, where the uncertainties are 1σ , defined such that 16% of the probability distribution is both above and below the error interval. If we omit object 7, then we have $\langle v^2 \rangle^{1/2} = 10.4^{+16.9}_{-1.2} \text{ km s}^{-1}$, and if we omit both 6 and 7, $\langle v^2 \rangle^{1/2} = 6.9^{+19.7}_{-0.4} \text{ km s}^{-1}$. For these three cases, in the same order, we are confident with 95% certainty that the dispersion is greater than 6.5, 7.3, and 4.7 km s^{-1} .

Assuming five genuine members with a true dispersion of 9.3 km s^{-1} , the systemic heliocentric velocity of UMajor is $-52.5 \pm 4.3 \text{ km s}^{-1}$.

4. MASS TO LIGHT RATIO

The central mass to light ratio of a stellar system with an isotropic orbital distribution and constant M/L may be expressed as $(M/L)_0 = 9 \eta \langle v^2 \rangle / (2\pi G r_{\text{hb}} S_0)$, where η is a luminosity distribution dependent factor always close to 1, r_{hb} is the half-light radius, and S_0 is the surface brightness (Richstone & Tremaine 1986). Using UMajor's approximate published total luminosity $M_V = -6.75$ ($L_V \sim 4 \times 10^4 L_\odot$) and half-light radius $r_{1/2} = 250 \text{ pc}$ (Willman et al. 2005), we obtain a central surface brightness of $S_0 = 0.11 L_\odot \text{ pc}^{-2}$. Assuming that $\langle v^2 \rangle^{1/2} = 9.3$, one then obtains $(M/L)_0 \approx 550 M_\odot / L_\odot$, as one might expect for a system that is nearly an order of magnitude fainter than known dark-matter dominated dSphs, but has the same dispersion. The central density is then $\rho_0 = 0.18 M_\odot \text{ pc}^{-3} = 6.8 \text{ GeV cm}^{-3}$ using the above equation, or $\rho_0 = 0.22 M_\odot \text{ pc}^{-3}$ using the formula $\rho_0 = 166 \langle v^2 \rangle / r_{\text{core}}$ (Mateo 1998). Despite UMajor's extreme M/L , its density is within a factor of two of that of several other dSphs (Mateo 1998). Several important caveats apply, however. First, we have identified the half-light radius with the half-brightness radius, when in truth they differ. If UMajor has a Plummer profile, for example, then $r_{\text{hb}} = 0.64 r_{1/2}$, and we underestimate M/L substantially. Next, the above formula for ρ_0/I_0 strictly applies only to systems with constant M/L , whereas a dwarf with a halo has varying M/L , a fact that is often ignored when computing M/L using core fitting. Thus the above values of $(M/L)_0$ and ρ_0 are intrinsically imprecise, and are useful primarily as relative values to compare with other dSphs having similarly computed qualities.

An alternative approach to computing the mass of the system is the projected mass estimator (Bahcall & Tremaine 1981; Heisler, Tremaine, & Bahcall 1985; Evans

et al. 2003), whereby the mass of a system with measured radial velocities v_i at projected radii R_i is given by $M = C/(GN) \times \sum_{i=1}^N v_i^2 R_i$, where C is a constant depending on the mass and light distributions of the system. For our five-member UMajor data, this becomes $M = C 1.5 \times 10^6 M_\odot$. Evans et al. (2003) compute C for the general case of measuring the mass between radii $r_{\min}$ to $r_{\max}$ for a $\rho \propto r^{-\gamma}$ tracer population in a $\phi \propto r^{-\alpha}$ potential. Taking $r_{\min} = 100$ pc from our innermost star and assuming that all the stars are within the half-light radius so that $r_{\max} = 250$ pc, and assuming that $\gamma = 3$, with a halo having a flat rotation curve ($\alpha = 0$), we obtain $C \approx 6$, again giving $M \approx 10^7 M_\odot$ and $M/L \sim 500$ inside the region observed. The mass varies from $5 \times 10^6 M_\odot$ to $2 \times 10^7 M_\odot$ as γ ranges from 2 to 6.

We further note that dSphs for which radially extended data exists have a mean M/L within the apparent stellar cutoff or “tidal” radius that is nearly an order of magnitude larger than the core-fit central M/L (e.g. compare Mateo (1998) with Kleyna et al. (2001), Wilkinson et al. (2004), and Kleyna et al. (2004)). If we accept the CDM result that halos are similar in structure, then UMajor might have a global $M/L \sim 3000 M_\odot/L_\odot$.

5. DISCUSSION

Our stellar velocity measurements show that UMajor is a nearly dark, low-mass halo, almost an order of magnitude less luminous than previously known dSph galaxies of similar mass. We obtain a central mass-to-light ratio $M/L \sim 500 M_\odot/L_\odot$; if this extrapolates to larger radii like other dSphs, UMajor may have a mean $M/L \sim 3000 M_\odot/L_\odot$. Hence, UMajor may be the most dark matter dominated galaxy yet discovered. For comparison, the recently discovered $10^{11} M_\odot$ VIRGOHI 21 HI source has $M/L > 500 M_\odot/L_\odot$ (Minchin et al. 2005), and the extreme dark spiral NGC 2915 has $M/L \approx 80$ (Meurer et al. 1996).

The CDM structure formation paradigm predicts that dark matter clumps into cusped halos characterized by a single parameter, the central density, with the smallest and least massive halos being the densest (Navarro, Frenk, & White 1997). Problematically, CDM also predicts an order of magnitude more small Galactic satellite halos than are actually observed (e.g. Klypin et al. 1999), but it is unclear whether the theory is at fault, or whether the missing halos are too dark to be observed.

High Velocity Clouds (HVCs) have been suggested as candidates for the missing CDM dwarf halos (Blitz et al. 1999). This suggestion remains controversial, especially as the high velocity clouds may themselves be divided into compact and diffuse subclasses with possibly different origins (e.g. de Heij, Braun, & Burton 2002). There is no clear distance determinant for the HVCs around the Milky Way, and, for a long time, there was no clear consensus as to whether the HVCs were Galactic or extra-Galactic. This ambiguity may have been resolved by the detection of a faint circumgalactic HVC population around the Andromeda galaxy by Thilker et al. (2004), who argue that the available data are consistent with formation mechanisms via cooling flows, or with tidal debris from mergers, or with gaseous counterparts of the missing CDM haloes.

Accordingly, UMajor may represent the best candidate

for a “missing” CDM halo. Its existence raises several interesting questions.

Why do the dSphs have a similar velocity dispersion, a similar central density and, presumably, a similar mass, as noted by Mateo et al. (1998)? It is curious that Draco, UMi, and UMajor, the three lowest luminosity Galactic dSphs, all have the same dispersion, about $\langle v^2 \rangle^{1/2} = 10 \text{ km s}^{-1}$, and And IX, the second faintest known dSph, has $\langle v^2 \rangle^{1/2} = 6.8_{-2.0}^{+3.0}$ (Chapman et al. 2005).

Is there a minimum halo size in which stars can form, or a minimum clustering scale for the dark matter? There was no survivability reason why UMajor could not have been much less massive. For instance, the mass required to bind a 500 pc dwarf halo against tidal disruption by a $10^{12} M_\odot$ Galaxy at UMajor’s distance of 100 kpc is estimated by equating the dwarf and Galactic densities, so that $M_{\text{bind}} \sim 10^{12} M_\odot \times (500 \text{ pc}/100 \text{ kpc})^3 \sim 10^5 M_\odot$, or about two orders of magnitude less massive than UMajor’s central region, and perhaps three orders of magnitude less massive than its global mass.

Kormendy & Freeman (2004) have compiled scaling relations among dwarf galaxies and more massive ellipticals, relating M_B , ρ_0 , r_{core} , and $\langle v^2 \rangle^{1/2}$. We can locate UMajor in this ensemble, using the values $\rho_0 = 0.18 M_\odot \text{ pc}^{-3}$, $\langle v^2 \rangle^{1/2} = 9.3 \text{ km s}^{-1}$, $r_{\text{core}} \sim r_{1/2} \sim 250 \text{ pc}$, and $M_B = M_V + 0.9 = -5.65$, where $M_V = -6.75$ has been converted to B band using the typical $B - V \approx 0.9$ color of a K-giant. From Figures 2 and 4 of Kormendy & Freeman (2004), we find, unsurprisingly, that UMajor is typical of the dSphs in all respects but luminosity. All but two of the previously known dSphs and dIrrs fall onto a tight Faber-Jackson relation between velocity dispersion and luminosity, albeit with a systematic offset relative to larger ellipticals. UMi and Draco are the exceptions, being about three magnitudes under-luminous for their mass. To agree with the general trend, UMajor would need to have a dispersion of $< 3 \text{ km s}^{-1}$, or would have to be about five magnitudes brighter.

It has been suggested that the dSphs’ extreme M/L results from the expulsion of baryons from their shallow potential well by supernovae (Larson 1974; Dekel & Silk 1986). Because the potential wells of the dSphs seem comparable, UMajor must have been intrinsically baryon poor to begin with, or else the baryon expulsion efficiency must have varied by one or two orders of magnitude among dSphs in order to produce their very different luminosities. UMajor demonstrates that the luminosity scatter at the low mass end of the galaxy distribution is very large. It appears likely that more dark and massive dwarfs are lurking in the vicinity of the Galaxy. Detections of more candidate objects are urgently needed to check whether the number and properties of the population are truly consistent with the missing dark matter haloes of the simulations.

The data presented herein were obtained at the W.M.

Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck

Foundation.

JTK gratefully acknowledges the support of the Beatrice Watson Parrent Fellowship. MIW acknowledges the support of PPARC.

REFERENCES

- Abazajian, K., Adelman, J., Agueros, M., et al. 2005, *AJ*, 129, 1755
 Armandroff, T. E., & Da Costa, G. S., 1991, *AJ*, 101, 1329
 Bahcall, J., & Tremaine, S., 1981, *ApJ*, 244, 805
 Blitz, L., Spergel, D. N., Teuben, P.J., Hartmann, D., & Burton, W.B., 1999, *ApJ*, 514, 818
 Chapman, S. C., Ibata, R., Lewis, G. R., Ferguson, A. M. N., Irwin, M., McConnachie, A., & Tanvir, N., 2005, accepted for publication in *ApJL* (astro-ph/0506103)
 Dekel, A., & Silk, J., 1986, *ApJ*, 303, 39
 Evans, N. W., Wilkinson, M. I., Perrett, K. M., & Bridges, T. J. 2003, *ApJ*, 583, 752
 Heisler, J., Tremaine, S., & Bahcall, J. N. 1985, *ApJ*, 298, 8
 de Heij V., Braun R., Burton W.B., 2002, *AA*, 392, 417
 Kleyna, J. T., Wilkinson, M. I., Evans, N. W., & Gilmore, G. 2001, *ApJL*, 563, L115
 Kleyna J. T., Wilkinson M. I., Evans N.W., Gilmore G., 2004, *MNRAS*, 354, L66
 Klypin, A., Kravtsov, A. V., Valenzuela, O., & Prada, F. 1999, *ApJ*, 522, 82
 Kormendy, J., & Freeman, K. C. 2004, *IAU Symposium*, 220, 377
 Larson, R.B., 1974, *MNRAS*, 169, 229
 Mateo M., 1998, *ARA&A*, 36, 435
 Mateo, M., Olszewski, E. W., Vogt, S. S., & Keane, M. J. 1998, *AJ*, 116, 2315
 Meurer, G. R., Carignan, C., Beaulieu, S. F., & Freeman, K. C. 1996, *AJ*, 111, 1551
 Minchin, R., et al. 2005, *ApJ*, 622, L21
 Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, *ApJ*, 490, 493
 Richstone, D.O., & Tremaine, S., 1986, *AJ*, 92, 72
 Thilker, D. A., Braun, R., Waltherbos, R. A. M., Corbelli, E., Lockman, F. J., Murphy, E., & Maddalena, R. 2004, *ApJ*, 601, L39
 Wilkinson, M. I., Kleyna, J. T., Evans, N. W., Gilmore, G. F., Irwin, M. J., & Grebel, E. K. 2004, *ApJ*, 611, L21
 Willman, B., et al., 2005, *ApJL*, submitted, (astro-ph/0503552)
 Vogt, S. S., et al. 1994, *Proc. SPIE*, 2198, 362
 Zucker, D. B., et al. 2004, *ApJ*, 612, L121




FIG. 1.— SDSS color-magnitude diagram for stars in the inner 6' of Ursa Major, with clearly visible giant and horizontal branches. The filled circles represent giant branch candidates observed in this work.




FIG. 2.— Velocities of Ursa Major candidate objects and σ_{scat} uncertainties. Stars 3 and 4 (near top) appear to be non-members, with velocities displaced about 60 km s⁻¹ from the others.




FIG. 3.— Equivalent widths of the three Calcium triplet lines. The reddest two lines of objects 3 and 4, which are apparent non-members on the basis of velocity, also have a significantly larger equivalent width than the other stars. The equivalent widths for object 7 are more uncertain than the error bars indicate because the absorption lines were almost invisible to the eye, and the fits may be entirely spurious. No fit could be obtained for the reddest line of object 7.

TABLE 1
POSITIONS, MAGNITUDES, AND VELOCITIES OF UMAJOR CANDIDATE STARS

Object	RA (J2000)	Dec	g	i	v (km s^{-1})	σ_{IRAF} (km s^{-1})	σ_{scat} (km s^{-1})
1	10 ^h 35 ^m 28 ^s .51	+51°57′00″.9	18.64	17.45	-54.87	1.00	1.30
2	10 ^h 34 ^m 30 ^s .51	+51°57′07″.0	18.88	17.75	-50.80	0.96	1.09
3	10 ^h 35 ^m 15 ^s .87	+51°59′32″.0	18.93	17.82	15.12	1.45	0.94
4	10 ^h 34 ^m 52 ^s .44	+51°57′02″.2	19.08	17.89	8.21	1.50	0.40
5	10 ^h 34 ^m 52 ^s .05	+51°58′28″.3	19.29	18.25	-64.22	1.90	0.65
6	10 ^h 34 ^m 42 ^s .36	+51°58′06″.1	19.45	18.37	-38.95	1.80	1.25
7	10 ^h 35 ^m 17 ^s .23	+51°55′33″.7	19.48	18.38	-53.53	5.00	5.15

Note. — First column is an object identification number. Columns 2 and 3 are the position in J2000 coordinates. Columns 4 and 5 are the SDSS g and i magnitudes. Columns 6, 7, and 8 are v , the measured heliocentric velocity; σ_{IRAF} , the uncertainty obtained from the rescaled IRAF FXCOR velocity error; and σ_{scat} , the uncertainty obtained from the velocity scatter among the different absorption lines.